

LA-UR-15-20565

Approved for public release; distribution is unlimited.

Title: Sign problem in Z-coefficient for particle emission angular distributions

Author(s): Kawano, Toshihiko

Intended for: communication with scientists outside LANL

Issued: 2015-01-29

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Sign problem in Z -coefficient for particle emission angular distributions

T. Kawano*

Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

(Dated: January 28, 2015)

* kawano@lanl.gov

There are two definitions of Blatt-Biedenharn's Z -coefficient, the one in the original paper [1]

$$Z(l_1 j_1 l_2 j_2; sL) = i^{-l_1+l_2+L} \hat{l}_1 \hat{l}_2 \hat{j}_1 \hat{j}_2 \langle l_1 l_2 00 | L0 \rangle W(l_1 j_1 l_2 j_2; sL), \quad (1)$$

and another one by Lane and Thomas [2]

$$\bar{Z}(l_1 j_1 l_2 j_2; sL) = \hat{l}_1 \hat{l}_2 \hat{j}_1 \hat{j}_2 \langle l_1 l_2 00 | L0 \rangle W(l_1 j_1 l_2 j_2; sL), \quad (2)$$

where the difference is the phase factor $i^{-l_1+l_2+L}$. We follow the notation by Lane and Thomas.

The angular distribution of shape elastic scattering (for neutrons) is given by the scattering amplitudes $A(\theta)$ and $B(\theta)$ at a given center-of-mass angle θ ;

$$A(\theta) = \frac{i}{2k} \sum_l \{ (l+1)(1 - S_{l,l+1/2}) + l(1 - S_{l,l-1/2}) \} P_l(\cos \theta), \quad (3)$$

$$B(\theta) = \frac{1}{2k} \sum_l \{ S_{l,l-1/2} - S_{l,l+1/2} \} P_l^1(\cos \theta), \quad (4)$$

$$\left(\frac{d\sigma}{d\Omega} \right)^{\text{SE}} = |A(\theta)|^2 + |B(\theta)|^2, \quad (5)$$

where S_{lj} is the optical model S -matrix element, $P_l(\cos \theta)$ is the Legendre function, and $P_l^1(\cos \theta)$ is the associated Legendre function. The same angular distribution can be expanded by the Legendre function as

$$\left(\frac{d\sigma}{d\Omega} \right)^{\text{SE}} = \sum_L B_L P_L(\cos \theta), \quad (6)$$

and the coefficients are given as

$$B_L = \frac{1}{8k^2} \sum_{l_1 j_1 l_2 j_2} \left[\bar{Z}(l_1 j_1 l_2 j_2; \frac{1}{2}L) \right]^2 \Re \{ (1 - S_{l_1 j_1})(1 - S_{l_2 j_2})^* \}. \quad (7)$$

This expression is independent of the target spin I_A . Since the \bar{Z} -coefficient is squared, the phase factor of $Z(l_1 j_1 l_2 j_2; 1/2L)$ does not matter. In this expansion, $4\pi B_0$ is equal to the angle-integrated elastic scattering cross section. To confirm the equality of Eq. (5) and Eq. (6) we calculated the shape elastic scattering angular distribution for ^{152}Gd at 20 MeV, with both equations, shown in Fig. 1. Koning-Delaroche global optical potential was used.

The differential cross section for the compound reaction is written in the same way as the shape elastic scattering;

$$\left(\frac{d\sigma}{d\Omega} \right)_{ab} = \sum_L B_L P_L(\cos \theta_b), \quad (8)$$

in which a particle a with the spin i_a collides with the target nucleus A with the spin I_A , forming a compound state J , then decays into a channel where a particle b with the spin i_b is emitted leaving the residual nucleus B with the spin I_B . Parity is not written explicitly, but it must be conserved at each spin coupling.

The B_L coefficient is given by Moldauer's statistical theory as

$$B_L = \frac{1}{4k^2} \frac{(-)^{I_B-I_A+i_b-i_a}}{(2i_a+1)(2I_A+1)} \frac{1}{N} \sum_J (2J+1)^2 \sum_{l_a j_a} \sum_{l_b j_b} W_{ab} \{ X_{l_a j_a}(E_a) X_{l_b j_b}(E_b) + \delta_{I_A I_B} \delta_{E_a E_b} Y_{l_a j_a, l_b j_b}(E_a, E_b) \}, \quad (9)$$

where W_{ab} is the width fluctuation factor, and

$$X_{lj}(E) = \bar{Z}(lj; iL) W(jJjJ; IL) T_{lj}(E), \quad (10)$$

$$Y_{l_a j_a, l_b j_b}(E_a, E_b) = (1 - \delta_{l_a l_b})(1 - \delta_{j_a j_b}) \{ \bar{Z}(l_a j_a l_b j_b; i_a L) W(J j_a J j_b; I_A L) \}^2 T_{l_a j_a}(E_a) T_{l_b j_b}(E_b) \quad (11)$$

and the normalization N is given by integrating/summing all possible decay channels.

$$N = \sum \int T_{lj}(E) dE. \quad (12)$$

For the Hauser-Feshbach theory, $W_{ab} = 1$ and $Y_{l_a j_a, l_b j_b}(E_a, E_b) = 0$.




FIG. 1. Calculated shape elastic scattering angular distribution of ^{152}Gd at 20 MeV. The symbols are from the scattering amplitudes at each angle, and the curve is from the Legendre coefficients.

The term that includes $X_{l_a j_a}(E_a)X_{l_b j_b}(E_b)$ has the factor of $\bar{Z}(l_a j_a l_a j_a; i_a L)\bar{Z}(l_b j_b l_b j_b; i_b L)$. When the Z -coefficients are given by the original definition,

$$\begin{aligned} Z(l_a j_a l_a j_a; i_a L)Z(l_b j_b l_b j_b; i_b L) &= i^{2L}(2l_a + 1)(2j_a + 1)(2l_b + 1)(2j_b + 1) \\ &\times \langle l_a l_a 00 | L0 \rangle \langle l_b l_b 00 | L0 \rangle \\ &\times W(l_a j_a l_a j_a; i_a L)W(l_b j_b l_b j_b; i_b L). \end{aligned} \quad (13)$$

Since L is always even for the compound reaction, i^{2L} becomes just one. Therefore the Legendre coefficient B_L will be identical for the both definition, Z and \bar{Z} (note that Z -coefficient in Y is squared). As an example, the neutron elastic scattering angular distributions for 2-MeV neutron induced reaction on ^{58}Ni are shown in Fig. 2, where the compound elastic scattering is 90-degree symmetric by definition.

[1] J.M. Blatt, L.C. Biedenharn, Rev Mod Phys. **24**, 258 (1952).

[2] A. M. Lane, R. G. Thomas, Rev. Mod. Phys. **30**, 257 (1958).




FIG. 2. Calculated shape and compound elastic scattering angular distributions for ^{58}Ni at 2 MeV.